

# TARO-FINGOLIMOD

## PREGNANCY-SPECIFIC PATIENT REMINDER CARD

- TARO-FINGOLIMOD is contraindicated during pregnancy and in women of childbearing potential not using effective contraception.
- Doctors will provide counselling before treatment initiation and regularly thereafter regarding the teratogenic risk of TARO-FINGOLIMOD and required actions to minimize this risk.
- Patients must use effective contraception while taking TARO-FINGOLIMOD.
- A pregnancy test must be carried out and negative results verified by the doctor before starting treatment. It must be repeated at suitable intervals.
- Patients will be informed by their doctor of the need for effective contraception while on treatment and for two months after discontinuation.

# TARO-FINGOLIMOD

## PREGNANCY-SPECIFIC PATIENT REMINDER CARD

- These are some sites for information on birth control options:
  - [sexandu.ca](http://sexandu.ca)
  - [plannedparenthood.org](http://plannedparenthood.org)

# TARO-FINGOLIMOD

## PREGNANCY-SPECIFIC PATIENT REMINDER CARD

- Doctors will provide counselling in the event of pregnancy and evaluation of the outcome of any pregnancy.
- While on treatment, women must not become pregnant. If a woman becomes pregnant or wants to become pregnant, TARO-FINGOLIMOD must be discontinued.
- Patients should inform their doctor immediately if there is worsening of multiple sclerosis after stopping treatment with TARO-FINGOLIMOD.
- TARO-FINGOLIMOD is secreted in breast milk. It is not recommended to breast feed an infant when taking this medication.

# TARO-FINGOLIMOD

## PREGNANCY-SPECIFIC PATIENT REMINDER CARD

### TARO-FINGOLIMOD and Male Reproduction

Available information does not suggest that TARO-FINGOLIMOD would be toxic to an unborn child when taken by the father.

- However, as a precaution, some clinicians will advise male patients to stop TARO-FINGOLIMOD at least three months before considering trying to father a child.

