HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use FENOFIBRATE TABLETS safely and effectively. See full prescribing information for FENOFIBRATE TABLETS.

FENOFIBRATE tablet, for oral use

Initial U.S. Approval: 1993 ------RECENT MAJOR CHANGES-

Warnings and Precautions, Hepatotoxicity (5.2) 03/2021 ----- INDICATIONS AND USAGE --

Fenofibrate tablets are a peroxisome proliferator-activated receptor (PPAR) alpha agonist indicated as an adjunct to diet: • To reduce elevated LDL-C, Total-C, TG and Apo B, and to increase HDL-C in adult activate with network here there are network deviced with advection (11)

- patients with primary hypercholesterolemia or mixed dyslipidemia (1.1). For treatment of adult patients with severe hypertriglyceridemia (1.2). Limitations of Use: Fenofibrate was not shown to reduce coronary heart disease morbidity and mortality in patients with type 2 diabetes mellitus (5.1).
- -- DOSAGE AND ADMINISTRATION --

Primary hypercholesterolemia or mixed dyslipidemia: Initial dose of 145 mg once

- daily (2.2). Severe hypertriglyceridemia: Initial dose of 48 to 145 mg once daily. Maximum dose is 145 mg (2.3).
- Renally impaired patients: Initial dose of 48 mg once daily (2.4), Geriatric patients: Select the dose on the basis of renal function (2.5). Maybe taken without regard to meals (2.1).
- -----DOSAGE FORMS AND STRENGTHS --

Oral Tablets: 48 mg and 145 mg (3)

----- CONTRAINDICATIONS -

- Severe renal dysfunction, including dialysis patients (4, 8.6, 12.3). Active liver disease (4, 5.2). Gallbladder disease (4, 5.5).
- Known hypersensitivity to fenofibrate (4).
- Nursing mothers (4, 8.2)

---- WARNINGS AND PRECAUTIONS ---

uced liver injury, including liver transplantation Hepatotoxicity: Serious drug-indu

FULL PRESCRIBING INFORMATION: CONTENTS

1 INDICATIONS AND USAGE Primary Hypercholesterolemia or Mixed Dyslipidemia

Severe Hypertriglyceridemia ns of Us

2 DOSAGE AND ADMINISTRATION

- General Considerations Primary Hypercholesterolemia or Mixed Dyslipidemia
- 2.2 Severe Hypertriglyceridemia 2.3
- 2.4 Impaired Renal Function
- 2.5 Geriatric Patient

3 DOSAGE FORMS AND STRENGTHS

- CONTRAINDICATIONS
- WARNINGS AND PRECAUTIONS
- Mortality and Coronary Heart Disease Morbidity
- Henatotoxicity 52 5.2 Hepatotoxicity
 5.3 Myopathy and Rhabdomyolysis
 5.4 Serum Creatinine
 5.5 Cholelithiasis
 5.6 Coumarin Anticoagulants
 5.7 Pancreatitis
 6.8 Heneral Market Alexandre

- 5.7 Fancteauus
 5.8 Hematologic Changes
 5.9 Hypersensitivity Reactions
 5.10 Venothromboembolic Disease
 5.11 Paradoxical Decreases in HDL Cholesterol Levels
- 6 ADVERSE REACTIONS Clinical Trials Experience

6.2 Postmarketing Experience

FULL PRESCRIBING INFORMATION **1 INDICATIONS AND USAGE**

1.1 Primary Hypercholesterolemia or Mixed Dyslipidemia

Fenofibrate tablets are indicated as adjunctive therapy to diet to reduce elevated lowdensity lipoprotein cholesterol (LDL-C), total cholesterol (Total-C), Triglycerides and apolipoprotein B (Apo B), and to increase high-density lipoprotein cholesterol (HDL-C) in adult patients with primary hypercholesterolemia or mixed dyslipidemia. 1.2 Severe Hypertriglyceridemia

Fenofibrate tablets are also indicated as adjunctive therapy to diet for treatment of adult patients with severe hypertriglyceridemia. Improving glycemic control in diabetic patients showing fasting chylomicronemia will usually obviate the need for pharmacologic intervention

Markedly elevated levels of serum triglycerides (e.g. > 2,000 mg/dL) may increase the risk of developing pancreatitis. The effect of fenofibrate therapy on reducing this risk has not been adequately studied.

1.3 Important Limitations of Use

Fenofibrate at a dose equivalent to 145 mg of fenofibrate tablets was not shown to reduce coronary heart disease morbidity and mortality in a large, randomized controlled trial of patients with type 2 diabetes mellitus [see Warnings and Precautions (5.1)]. 2 DOSAGE AND ADMINISTRATION

2.1 General Considerations

Patients should be placed on an appropriate lipid-lowering diet before receiving fanofibrate tablets, and should continue this diet during treatment with fenofibrate tablets. Fenofibrate tablets can be given without regard to meals.

The initial treatment for dyslipidemia is dietary therapy specific for the type of lipoprotein The initial readiness to dysplotenia is breatly the ady specific of the type of npoprotein abnormality. Excess body weight and excess alcoholic intake may be important factors in hypertriglyceridemia and should be addressed prior to any drug therapy. Physical exercise can be an important ancillary measure. Diseases contributory to hyperlipidemia, such as hypothyroidism or diabetes mellitus should be looked for and adequately treated. Estrogen therapy, thiazide diuretics and beta-blockers, are sometimes associated with massive rises in plasma triglycerides, especially in subjects with familial hypertriglyceridemia. In such cases, discontinuation of the specific etiologic agent may viate the need for specific drug therapy of hypertriglyceridemia.

Lipid levels should be monitored periodically and consideration should be given to reducing the dosage of fenofibrate tablets if lipid levels fall significantly below the

Therapy should be withdrawn in patients who do not have an adequate response after two months of treatment with the maximum recommended dose of 145 mg once daily. 2.2 Primary Hypercholesterolemia or Mixed Dyslipidemia

The initial dose of fenofibrate tablets is 145 mg once daily

2.3 Severe Hypertriglyceridemia

and death, has been reported with fenofibrate. Monitor patient's liver function, including serum ALT, AST, and total bilirubin, at baseline and periodically for the duration of therapy. Discontinue if signs or symptoms of liver injury develop or if elevated enzyme levels persist (5.2).

- Myopathy and rhabdomyolysis: Have been reported in patients taking fenofibrate. Risks are increased during co-administration with a statin (with a significantly higher rate observed for gemfibrozil), particularly in elderly patients and patients
- with diabetes, renal failure, or hypothyroidism (5.3) Serum Creatinine: Fenofibrate can reversibly increase serum creatinine levels
- Security Detailines renormate can reversibly inclease security total mereters (5.4). Monitor renal function periodically in patients with renal impairment (8.6). *Cholelithiasis*: Fenofibrate increases cholesterol excretion into the bile, leading to risk of cholelithiasis. If cholelithiasis is suspected, gallbladder studies are indicated (5.5).
- Coumarin anticoagulants: Use caution in concomitant treatment with oral coumarin
- Cournari anticcaguiants: Use caution in concomitant treatment with oral cournarin anticcaguiants. Adjust the dosage of cournarin anticcaguiant to maintain the prothrombin time/INR at the desired level to prevent bleeding complications (5.6). *Hypersensitivity Reactions:* Acute hypersensitivity reactions, including severe cutaneous adverse drug reactions have been reported postmarketing. Some cases were life-threatening and required emergency treatment. Discontinue

fenofibrate and treat patients appropriately if reactions occur (5.9).

----- ADVERSE REACTIONS ---Adverse reactions > 2% and at least 1% greater than placebo: Abnormal liver tests, increased AST, increased ALT, increased CPK, and rhinitis (6). To report SUSPECTED ADVERSE REACTIONS, contact Sum Pharmaceutical Industries Inc. at 1-800-406-7984 or FDA at 1-800-FDA-1088 or www.ida.gov/medwatch

----- DRUG INTERACTIONS ----

- Coumarin anticoagulants: (7.1
- Immunosuppressants: (7.2). Bile acid resins: (7.3).

--- USE IN SPECIFIC POPULATIONS Geriatric Use: Determine dose selection based on renal function (8.5)

Renal Impairment: Avoid use in severe renal impairment patients. Dose reduction is required in mild to moderate renal impairment patients (8.6)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 07/2021

7 DRUG INTERACTIONS
7.1 Coumarin Anticoagulants
7.2 Immunosuppressants
7.3 Bile Acid Binding Resins
7.4 Colchicine
8 USE IN SPECIFIC POPULATIONS
8.1 Pregnancy
8.2 Lactation
8.4 Pediatric Use
8.5 Geriatric Use
8.6 Renal Impairment
8.7 Hepatic Impairment
10 OVERDOSAGE
11 DESCRIPTION
12 CLINICAL PHARMACOLOGY
12.1 Mechanism of Action
12.2 Pharmacodynamics
12.3 Pharmacokinetics
13 NONCLINICAL TOXICOLOGY
13.1 Carcinogenesis and Mutagenesis and Impairment of Fertility
14 CLINICAL STUDIES
14.1 Primary Hypercholesterolemia (Heterozygous Familial and No

nd Nonfamilial) and Mixed Dyslipidemia

14.2 Severe Hypertriglyceridemia 16 HOW SUPPLIED/STORAGE AND HANDLING **17 PATIENT COUNSELING INFORMATION**

Sections or subsections omitted from the full prescribing information are not listed.

to patient response, and should be adjusted if necessary following repeat lipid determinations at 4 to 8 week intervals. The maximum dose is 145 mg once daily. 2.4 Impaired Renal Function

Treatment with fenofibrate tablets should be initiated at a dose of 48 mg per day in patients having mild to moderately impaired renal function, and increased only after evaluation of the effects on renal function and lipid levels at this dose. The use of fenofibrate tablets should be avoided in patients with severe renal impairment [see Use in Specific Populations (8.6) and Clinical Pharmacology (12.3)].

2.5 Geriatric Patients

Dose selection for the elderly should be made on the basis of renal function [see Use in Specific Populations (8.5)]

3 DOSAGE FORMS AND STRENGTHS

- 48 mg white to off-white, oval-shaped, biconvex, film-coated tablets debossed with 'RH37' on one side and other side plain.
- 145 mg white to off-white, oval-shaped, biconvex, film-coated tablets debossed with 'RH38' on one side and other side plain.

4 CONTRAINDICATIONS

Fenofibrate is contraindicated in:

- patients with severe renal impairment, including those receiving dialysis [see Clinical Pharmacology (12.3)].
- patients with active liver disease, including those with primary biliary cirrhosis and unexplained persistent liver function abnormalities [see Warnings and Precautions (5.2)].
- patients with preexisting gallbladder disease [see Warnings and Precautions (5.5)].
- nursing mothers [see Use in Specific Populations (8.2)].
- patients with known hypersensitivity to fenofibrate or fenofibric acid [see Warnings and Precautions (5.9)]. **5 WARNINGS AND PRECAUTIONS**

5.1 Mortality and Coronary Heart Disease Morbidity

The effect of fenofibrate on coronary heart disease morbidity and mortality and noncardiovascular mortality has not been established.

The Action to Control Cardiovascular Risk in Diabetes Lipid (ACCORD Lipid) trial was a randomized placebo-controlled study of 5518 patients with type 2 diabetes mellitus on background statin therapy treated with fenofibrate. The mean duration of follow-up was 4.7 years. Fenofibrate plus statin combination therapy showed a non-significant 8% relative risk reduction in the primary outcome of major adverse cardiovascula or relative has reduction in the primary outcome to map a worse catavasculat events (MACE), a composite of non-fatal myocardial infraction, non-fatal stroke, and cardiovascular disease death (hazard ratio [HR] 0.32, 95% CI 0.79 to 1.08) (p = 0.32) as compared to statim monotherapy. In a gender subgroup analysis, the hazard ratio for MACE in men receiving combination therapy versus statin monotherapy was 0.82 (95% CI 0.69 to 0.99), and the hazard ratio for MACE in women receiving combination

0.01). The clinical significance of this subgroup finding is unclear The Fenofibrate Intervention and Event Lowering in Diabetes (FIELD) study was a 5-year randomized, placebo-controlled study of 9795 patients with type 2 diabetes mellitus treated with fenofibrate. Fenofibrate demonstrated a non-significant 11% The interval of the interval is the interval of the interval

and white blood cell counts are recommended during the first 12 months of

Anaphylaxis and angioedema have been reported postmarketing with fenofibrate. In

some cases, reactions were life-threatening and required emergency treatment. If a patient develops signs or symptoms of an acute hypersensitivity reaction, advise them

Severe cutaneous adverse drug reactions (SCAR), including Stevens-Johnson

syndrome, toxic epidermal necrolysis, and Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS), have been reported postmarketing, occurring days to

weeks after initiation of fenofibrate. The cases of DRESS were associated with

vectors and initial and ini

In the FIELD trial, pulmonary embolus (PE) and deep vein thrombosis (DVT) were

observed at higher rates in the fenofibrate- than the placebo-treated group. Of

business and the second secon

In the Coronary Drug Project, a higher proportion of the clofibrate group experienced definite or suspected fatal or nonfatal pulmonary embolism or thrombophlebitis than the placebo group (5.2% vs. 3.3% at five years; p < 0.01).

There have been postmarketing and clinical trial reports of severe decreases in HDL cholesterol levels (as low as 2 mg/dL) occurring in diabetic and non-diabetic patients initiated on fibrate therapy. The decrease in HDL-C is mirrored by a decrease in apolipoprotein A1. This decrease has been reported to occur within 2 weeks to years after initiation of fibrate therapy. The HDL-C levels remain depressed until fibrate

therapy has been withdrawn; the response to withdrawal of fibrate therapy is rapid and

sustained. The clinical significance of this decrease in HDL-C is unknown. It is recommended that HDL-C levels be checked within the first few months after initiation of fibrate therapy. If a severely depressed HDL-C level is detected, fibrate therapy should be withdrawn, and the HDL-C level monitored until it has returned to baseline,

The following serious adverse reactions are described below and elsewhere in the

· Mortality and coronary heart disease morbidity [see Warnings and Precautions

Because clinical studies are conducted under widely varying conditions, adverse reaction

rates observed in the clinical studies of a drug cannot be directly compared to rates in the clinical studies of another drug and may not reflect the rates observed in practice.

Adverse events reported by 2% or more of patients treated with fenofibrate (and greater than placebo) during the double-blind, placebo-controlled trials, regardless of causality, are listed in Table 1 below. Adverse events led to discontinuation of treatment in 5% of patients treated with fenofibrate and in 3% treated with placebo.

Increases in liver function tests were the most frequent events, causing discontinuation of fenofibrate treatment in 1.6% of patients in double-blind trials.

Table 1. Adverse Reactions Reported by 2% or More of Patients Treated with Fenofibrate and Greater than Placebo During the Double-Blind,

Placebo-Controlled Trials

Urticaria was seen in 1.1% vs. 0%, and rash in 1.4% vs. 0.8% of fenofibrate and placebo patients respectively in controlled trials.

In a pooled analysis of 10 placebo-controlled trials, increases to > 3 times the upper limit of normal in ALT occurred in 5.3% of patients taking fenofibrate at doses equivalent to 66 mg to 145 mg fenofibrate daily versus 1.1% of patients treated with placebo [see Warnings and Precautions (5.2)]. In an 8-week study, the incidence of

ALT or AST elevations \geq 3 times the upper limit of normal was 13% in patients

receiving dosages equivalent to 96 mg to 145 mg fenofibrate daily and was 0% in those receiving dosages equivalent to 48 mg or less fenofibrate daily or placebo.

The following adverse reactions have been identified during postapproval use of

fenofibrate. Because these reactions are reported voluntarily from a population of

uncertain size, it is not always possible to reliably estimate their freque poyer establish a causal relationship to drug exposure: myalgia, rhabdomyolysis, pancreatitis, acute renal failure, muscle spasm, hepatitis, cirrhosis, increased total bilinubin, anemia, arthralgia, decreases in hemoglobin, decreases in hematocrit, white blood cell

decreases, asthenia, severely depressed HDL-cholesterol levels, and interstitial lung

disease. Photosensitivity reactions have occurred days to months after initiation; in

ne of these cases, patients reported a prior photosensitivity reaction to ketoprot

Potentiation of coumarin-type anticoagulant effects has been observed with

Fenofibrate

(N = 439)

4.6%

3.4%

3.2%

2.3%

2.1%

7.5%

3%

3.4%

6.2%

2.3%

Placebo

(N = 365)

4 4%

2.5%

2.7%

1.9%

1.4%

1.4%

1.6%

1.4%

0.5%

1.1%

Hypersensitivity reactions [see Warnings and Precautions (5.9)] Venothromboembolic disease [see Warnings and Precautions (5.10)]

Hepatotoxicity [see Warnings and Precautions (5.2)]

Pancreatitis [see Warnings and Precautions (5.7)]

5.11 Paradoxical Decreases in HDL Cholesterol Levels

and fibrate therapy should not be re-initiated.

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

to seek immediate medical attention and discontinue fenofibrate

fenofibrate administration

Acute Hypersensitivity

Delayed Hypersensitivity

0.022).

labeling:

(5.1)]

BODY SYSTEM

Adverse Reaction

Abdominal Pair

Back Pain

Headache

DIGESTIVE

ncreased AL

Increased CPK

ncreased AS

RESPIRATOR

Rhinitis

Respiratory Disorder

Nausea

BODY AS A WHOLE

METABOLIC AND NUTRITIONAL DISORDERS

quivalent to 145 mg fenofibrat

Abnormal Liver Function Tes

"Significantly different from Placebo.

Increases in Liver Enzymes

6.2 Postmarketing Experience

7 DRUG INTERACTIONS

7.1 Coumarin Anticoagulants

prolongation of the PT/INR.

5 10 Venothromboembolic Disease

5.9 Hypersensitivity Reactions

J. USP

FENOFIBRAT ЩS, TABL

respectively, with fenofibrate as compared to placebo. Because of chemical, pharmacological, and clinical similarities between fenofibrate tablets, clofibrate, and gemfibrozil, the adverse findings in 4 large randomized, placebo-controlled clinical studies with these other fibrate drugs may also apply to fenofibrate. In the Coronary Drug Project, a large study of post myocardial infarction of patients treated for 5 years with clofibrate, there was no difference in mortality seen between the clofibrate group and the placebo group. There was however, a difference in the rate of cholelithiasis and cholecystitis requiring surgery between the two groups (3% vs. 1.8%). In a study conducted by the World Health Organization (WHO), 5000 subjects without known coronary artery disease were treated with placebo or clofibrate for 5 years and followed for an additional one year. There was a statistically significant, higher age -adjusted all-cause mortality in the clofibrate group compared with the placebo group to be a calculate of the second seco

The Helsinki Heart Study was a large (n = 4081) study of middle-aged men without a history of coronary artery disease. Subjects received either placebo or gemfibrozil for 5 years, with a 3.5 year open extension afterward. Total mortality was numerically higher in the gemfibrozil randomization group but did not achieve statistica significance (p = 0.19, 95% confidence interval for relative risk G:P = 0.91 to 1.64) Although cancer deaths trended higher in the gemfbrozil group (p = 0.11), cancers (excluding basal cell carcinoma) were diagnosed with equal frequency in both study groups. Due to the limited size of the study, the relative risk of death from any cause was not shown to be different than that seen in the 9 year follow-up data from World Health Organization study (RR = 1.29).

A secondary prevention component of the Helsinki Heart Study enrolled middle-aged men excluded from the primary prevention study because of known or suspected coronary heart disease. Subjects received gemfibrozil or placebo for 5 years. Although cardiac deaths trended higher in the gemfibrozil group, this was not statistically significant (hazard ratio 2.2. 95% confidence interval; 0.94 to 5.05). The rate of significant (nazive tato 2.2, 50 contestion marked or 50 cost of 0.00) and the cost of th 5.2 Hepatotoxicity

Serious drug-induced liver injury (DILI), including liver transplantation and death,

have been reported postmarketing with fenofibrate. DILI has been reported within the first few weeks of treatment or after several months of therapy and in some cases has reversed with discontinuation of fenofibrate treatment. Patients with DILI have

experienced signs and symptoms including dark urine, abnormal stool, jaundice, malaise, abdominal pain, myalgia, weight loss, pruritus, and nausea. Many patients had concurrent elevations of total bilirubin, serum alanine transaminase (ALT), and

aspartate transaminase (AST). DILI has been characterized as hepatocellular, chronic

active, and cholestatic hepatitis, and cirrhosis has occurred in association with chronic

In clinical trials, fenofibrate at doses equivalent to 96 mg to 145 mg fenofibrate tablets daily has been associated with increases in serum AST or ALT. The incidence of increases in transaminases may be dose related [see Adverse Reactions (6.1)].

Fenofibrate is contraindicated in patients with active liver disease, including those with primary biliary cirrhosis and unexplained persistent liver function abnormalities

[see Contraindications (4)]. Monitor patient's liver function, including serum ALT,

AST, and total bilirubin, at baseline and periodically for the duration of therapy with

For an total billiobin, at baseline and periodically for the outation of therapy with fenofibrate. Discontinue fenofibrate if signs or symptoms of liver injury develop or if elevated enzyme levels persist (ALT or AST > 3 times the upper limit of normal, or if accompanied by elevation of bilirubin). Do not restart fenofibrate in these patients if

Fibrates increase the risk for myopathy and have been associated with rhabdomyolysis

The risk for serious muscle toxicity appears to be increased in elderly patients and in

The new of all reserves indicated with a second sec

Patients should be advised to report promptly unexplained muscle pain, tenderness or weakness, particularly if accompanied by malaise or fever. CPK levels should be assessed in patients reporting these symptoms, and fenofibrate therapy should be

discontinued if markedly elevated CPK levels occur or myopathy/myositis is suspected

Data from observational studies indicate that the risk for rhabdomyolysis is increased

when fibrates, in particular gemfibrozil, are co-administered with a statin. The combination should be avoided unless the benefit of further alterations in lipid levels

is likely to outweigh the increased risk of this drug combination [see Clinical

Cases of myopathy, including rhabdomyolysis, have been reported with fenofibrates

co-administered with colchicine, and caution should be exercised when prescribing

Elevations in serum creatinine have been reported in patients on fenofibrate. These elevations tend to return to baseline following discontinuation of fenofibrate. The clinical significance of these observations is unknown. Monitor renal function in

patients with renal impairment taking fenofibrate. Renal monitoring should also be

considered for patients taking fenofibrate at risk for renal insufficiency such as the

Fenofibrate, like clofibrate and gemfibrozil, may increase cholesterol excretion into the bile, leading to cholelithiasis. If cholelithiasis is suspected, gallbladder studies are

Construction of the constr

Pancreatitis has been reported in patients taking fenofibrate, gemfibrozil, and clofibrate. This occurrence may represent a failure of efficacy in patients with severe hypertriglyceridemia, a direct drug effect, or a secondary phenomenon mediated

hrough biliary tract stone or sludge formation with obstruction of the common bile

indicated. Fenofibrate therapy should be discontinued if gallstones are found.

nofibrate with colchicine [see Drug Interactions (7.4)].

or weakness, and/or marked elevations of creatine phosphokinase (CPK) levels.

there is no alternative explanation for the liver injury.

5.3 Myopathy and Rhabdomyolysis

active hepatitis.

or diagnosed.

cology (12.3)].

Iderly and patients with diabetes.

5.6 Coumarin Anticoagulants

5.4 Serum Creatinine

5.5 Cholelithiasis

(7.1)1

duct

5.7 Pancreatitis

5.8 Hematologic Changes

The initial dose is 48 to 145 mg per day. Dosage should be individualized according

therapy versus statin monotherapy was 1.38 (95% CI 0.98 to 1.94) (interaction p = Mild to moderate hemoglobin, hematocrit, and white blood cell decreases have been observed in patients following initiation of fenofibrate therapy. However, these levels stabilize during long-term administration. Thrombocytopenia and agranulocytosis Caution should be exercised when coumarin anticoagulants are given in conj have been reported in individuals treated with fenofibrate. Periodic monitoring of red with fenofibrate. The dosage of the anticoagulants should be reduced to maintain the

PI Size: 273 x 314 mm Folded Size : 35 x 35 mm Font Size : 6 Pt. Color : Black Track : A23/06/2021, A28/06/2021, A29/06/2021. A03/07/2021



SAP Code: 5221486 ITF Code : 05221486 Supersede : 5198365 PT/INR at the desired level to prevent bleeding complications. Frequent PT/INR determinations are advisable until it has been definitely determined that the PT/INR has stabilized [see Warnings and Precautions (5.6)].

7.2 Immunosuppressants

Immunosuppressants such as cyclosporine and tacrolimus can produce nephrotoxicity with decreases in creatinine clearance and rises in serum creatinine, and because renal excretion is the primary elimination route of fibrate drugs including fenofibrate, there is a risk that an interaction will lead to deterioration of renal function. The benefits and risks of using fenofibrate tablets with immunosuppressants and other potentially nephrotoxic agents should be carefully considered, and the lowest effective dose employed and renal function monitored.

7.3 Bile Acid Binding Resins

Since bile acid binding resins may bind other drugs given concurrently, patients should take fenofibrate at least 1 hour before or 4 to 6 hours after a bile acid binding resin to avoid impeding its absorption.

7.4 Colchicine

Cases of myopathy, including rhabdomyolysis, have been reported with fenofibrates co-administered with colchicine, and caution should be exercised when prescribing fenofibrate with colchicine.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Limited available data with fenofibrate use in pregnant women are insufficient to determine a drug associated risk of major birth defects, miscarriage or adverse maternal or fetal outcomes. In animal reproduction studies, no evidence of embryo-fetal toxicity was observed with oral administration of fenofibrate in rats and rabbits during organogenesis at doses less than or equivalent to the maximum recommended clinical dose of 145 mg daily, based on body surface area (mg/m²). Adverse reproductive outcomes occurred at higher doses in the presence of maternal toxicity (see Data). Fenofibrate should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

<u>Data</u> Animal Data

In pregnant rats given oral dietary doses of 14, 127, and 361 mg/kg/day from gestation day 6 to 15 during the period of organogenesis, no adverse developmental findings were observed at 14 mg/kg/day (less than the clinical exposure at the maximum recommended human dose [MRHD] of 300 mg fenofibrate daily, equivalent to 145 mg fenofibrate daily, based on body surface area comparisons). Increased fetal skeletal malformations were observed at maternally toxic doses (361 mg/kg/day, corresponding to 12 times the clinical exposure at the MRHD) that significantly suppressed maternal body weight gain.

In pregnant rabbits given oral gavage doses of 15, 150, and 300 mg/kg/day from gestation day 6 to 18 during the period of organogenesis and allowed to deliver, no adverse developmental findings were observed at 15 mg/kg/day (a dose that approximates the clinical exposure at the MRHD, based on body surface area comparisons). Aborted litters were observed at maternally toxic doses (\geq 150 mg/kg/day, corresponding to \geq 10 times the clinical exposure at the MRHD) that suppressed maternal body weight gain.

In pregnant rats given oral dietary doses of 15, 75, and 300 mg/kg/day from gestation day 15 through lactation day 21 (weaning), no adverse developmental effects were observed at 15 mg/kg/day (less than the clinical exposure at the MRHD, based on body surface area comparisons), despite maternal toxicity (decreased weight gain). Postimplantation loss was observed at 2.75 mg/kg/day (≥ 2 times the clinical exposure at the MRHD) in the presence of maternal toxicity (decreased weight gain). Decreased pup survival was noted at 300 mg/kg/day (10 times the clinical exposure at the MRHD), which was associated with decreased maternal body weight gain/maternal neglect. 8.21 action

Risk Summary

There is no available information on the presence of fenofibrate in human milk, effects of the drug on the breastfed infant, or the effects on milk production. Fenofibrate is present in the milk of rats, and is therefore likely to be present in human milk. Because of the potential for serious adverse reactions in breastfed infants, such as disruption of infant lipid metabolism, women should not breastfeed during treatment with fenofibrate and for 5 days after the final dose [see Contraindications (4)]. 8.4 Pediatric Use

8.4 Pediatric Use

Safety and effectiveness have not been established in pediatric patients

8.5 Geriatric Use

Fenofibric acid is known to be substantially excreted by the kidney, and the risk of adverse reactions to this drug may be greater in patients with impaired renal function. Fenofibric acid exposure is not influenced by age. Since delarly patients have a higher incidence of renal impairment, dose selection for the elderly should be made on the basis of renal function [*see Dosage and Administration* (2.5) and *Clinical Pharmacology* (12.3)]. Elderly patients with normal renal function should require no dose modifications. Consider monitoring renal function in elderly patients taking fenofibrate.

8.6 Renal Impairment

The use of fenofibrate should be avoided in patients who have severe renal impairment [see Contraindications (4)]. Dose reduction is required in patients with mild to moderate renal impairment [see Dosage and Administration (2.4) and Clinical Pharmacology (12.3)]. Monitoring renal function in patients with renal impairment is recommended.

8.7 Hepatic Impairment


The use of fenofibrate has not been evaluated in subjects with hepatic impairment [see Contraindications (4) and Clinical Pharmacology (12.3)].

10 OVERDOSAGE

There is no specific treatment for overdose with fenofibrate. General supportive care of the patient is indicated, including monitoring of vital signs and observation of clinical status, should an overdose occur. If indicated, elimination of unabsorbed drug should be achieved by emesis or gastric lavage; usual precautions should be observed to maintain the airway. Because fenofibric acid is highly bound to plasma proteins, hemodialviss should no tbe considered.

11 DESCRIPTION

Fenofibrate, USP is a lipid regulating agent available as tablets for oral administration. Each tablet contains 48 mg or 145 mg of fenofibrate, USP. The chemical name for fenofibrate, USP is 2-(4-chorobenzoyl) phenoxyl-2-methyl-propanoic acid, 1-methylethyl ester with the following structural formula:



The molecular formula is $C_{20}H_{21}O_4Cl$ and the molecular weight is 360.83; fenofibrate, USP is very soluble in methylene chloride, slightly soluble in alcohol and practically insoluble in water.

The melting point is 79-82° C. Fenofibrate, USP is a white or almost white crystalline powder.

Inactive Ingredients

Each film-coated tablet contains colloidal silicon dioxide, crospovidone, hypromellose (3 cps), microcrystalline cellulose, polyvinyl alcohol, pregelatinized starch, sucrose, sodium lauryl sulfate, sodium stearyl fumarate, soybean lecithin, talc, titanium dioxide, and xanthan gum. Meets USP Dissolution Test 1.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The active moiety of fenofibrate is fenofibric acid. The pharmacological effects of fenofibric acid in both animals and humans have been extensively studied through oral administration of fenofibrate.

The lipid-modifying effects of fenofibric acid seen in clinical practice have been explained *in vivo* in transgenic mice and *in vitro* in human hepatocyte cultures by the activation of peroxisome proliferator activated receptor (PPARe). Through this mechanism, fenofibrate increases lipolysis and elimination of triglyceride-rich particles from plasma by activating lipoprotein lipase and reducing production of apoprotein C-III (an inhibitor of lipoprotein lipase.

The resulting decrease in TG produces an alteration in the size and composition of LDL from small, dense particles (which are thought to be atherogenic due to their susceptibility to oxidation), to large buoyant particles. These larger particles have a greater affinity for cholesterol receptors and are catabolized rapidly. Activation of PPARc also induces an increase in the synthesis of apolipoproteins A-I, A-II and HDLcholesterol.

Fenofibrate also reduces serum uric acid levels in hyperuricemic and normal individuals by increasing the urinary excretion of uric acid.

12.2 Pharmacodynamics

A variety of clinical studies have demonstrated that elevated levels of total-C, LDL-C, and apo B, an LDL membrane complex, are associated with human atherosclerosis. Similarly, decreased levels of HDL-C and its transport complex, apolipoprotein A (apo A land apo AII) are associated with the development of atherosclerosis. Epidemiologic investigations have established that cardiovascular morbidity and mortality vary directly with the level of total-C, LDL-C, and TG, and inversely with the level of HDL-C. The independent effect of raising HDL-C or lowering triglycerides (TG) on the risk of cardiovascular morbidity and mortality has not been determined.

Fenofibric acid, the active metabolite of fenofibrate, produces reductions in total cholesterol, LDL cholesterol, apolipoprotein B, total triglycerides and triglyceride rich lipoprotein (VLDL) in treated patients. In addition, treatment with fenofibrate results in increases in high density lipoprotein (HDL) and apolipoproteins apoAl and apoAll. **12.3 Pharmacokinetics**

Plasma concentrations of fenofibric acid after administration of three 48 mg or one 145 mg tablets are equivalent under fed conditions to one 200 mg micronized fenofibrate capsule.

Fenofibrate is a pro-drug of the active chemical moiety fenofibric acid. Fenofibrate is converted by ester hydrolysis in the body to fenofibric acid which is the active constituent measurable in the circulation.

Absorption

The absolute bioavailability of fenofibrate cannot be determined as the compound is virtually insoluble in aqueous media suitable for injection. However, fenofibrate is well absorbed from the gastrointestinal tract. Following oral administration in healthy volunteers, approximately 60% of a single dose of radiolabelled fenofibrate appeared in urine, primarily as fenofibric acid and its glucuronate conjugate, and 25% was excreted in the feces. Peak plasma levels of fenofibric acid occur within 6 to 8 hours after administration.

Exposure to fenofibric acid in plasma, as measured by C_{max} and AUC, is not significantly different when a single 145 mg dose of fenofibrate is administered under fasting or nonfasting conditions. Distribution

Upon multiple dosing of fenofibrate, fenofibric acid steady state is achieved within 9 days. Plasma concentrations of fenofibric acid at steady state are approximately double of those following a single dose. Serum protein binding was approximately 99% in normal and hyperlipidemic subjects. Metabolism

Following oral administration, fenofibrate is rapidly hydrolyzed by esterases to the active metabolite, fenofibric acid; no unchanged fenofibrate is detected in plasma. Fenofibric acid is primarily conjugated with glucuronic acid and then excreted in urine. A small amount of fenofibric acid is reduced at the carbonyl moiety to a benzhydrol metabolite which is, in turn, conjugated with glucuronic acid and excreted in urine.

In vivo metabolism data indicate that neither fenofibrate nor fenofibric acid undergo oxidative metabolism (e.g., cytochrome P450) to a significant extent. Elimination

After absorption, fenofibrate is mainly excreted in the urine in the form of metabolites, primarily fenofibric acid and fenofibric acid glucuronide. After administration of radiolabelled fenofibrate, approximately 60% of the dose appeared in the urine and 25% was excreted in the feces.

Fenofibric acid is eliminated with a half-life of 20 hours, allowing once daily dosing. Special Populations Geriatrics

In elderly volunteers 77 to 87 years of age, the oral clearance of fenofibric acid following a single oral dose of fenofibrate was 1.2 L/h, which compares to 1.1 L/h in young adults. This indicates that a similar dosage regimen can be used in elderly with normal renal function, without increasing accumulation of the drug or metabolites [see Dosage and Administration (2.5) and Use in Specific Populations (8.5)]. Pediatrics

The pharmacokinetics of fenofibrate has not been studied in pediatric populations Gender

No pharmacokinetic difference between males and females has been observed for fenofibrate.

The influence of race on the pharmacokinetics of fenofibrate has not been studied, however fenofibrate is not metabolized by enzymes known for exhibiting inter-ethnic variability.

Renal Impairment

Race

The pharmacokinetics of fenofibric acid was examined in patients with mild, moderate, and severe renal impairment. Patients with severe renal impairment (estimated glomerular filtration rate [eGFR] < 30 mL/min/1.73 m²) showed 2.7-fold increase in exposure for fenofibric acid and increased accumulation of fenofibric acid during chronic dosing compared to that of healthy subjects. Patients with mild to moderate renal impairment (eGFR 30 to 59 mL/min/1.73 m²) had similar exposure but an increase in the hall-life for fenofibric acid compared to that of healthy subjects. Based on these findings, the use of fenofibrate should be avoided in patients having mild to moderate renal impairment and dose reduction is required in patients having mild to moderate renal impairment *(see Dosage and Administration (2.4)]*.

Hepatic Impairment No pharmacokinetic studies have been conducted in patients with hepatic impairment.

armacokineuc suules have been conducted in patients with hepatic impairment

Drug-drug Interactions

In vitro studies using human liver microsomes indicate that fenofibrate and fenofibric acid are not inhibitors of cytochrome (CYP) P450 isoforms CYP3A4, CYP2D6 CYP2E1, or CYP1A2. They are weak inhibitors of CYP2C8, CYP2C19 and CYP2A6 and mild-to-moderate inhibitors of CYP2C9 at therapeutic concentrations.

Table 2 describes the effects of co-administered drugs on fenofibric acid systemic exposure. Table 3 describes the effects of co-administered fenofibrate or fenofibric acid on other drugs.

Table 2. Effects of Co-Administered Drugs on Fenofibric Acid Systemic Exposure from Fenofibrate Administration

Co- Administered	ninistered Dosage Regimen of Dosage Regimen		Changes in	Fenofibric
Drug	Co-Administered Drug	of Fenofibrate	Acid Exposure	
			AUC	C _{max}
Lipid-lowering agents				
Atorvastatin	20 mg once daily for for 10 days	Fenofibrate 160 mg ¹ once daily for 10 days	↓2%	↓4%
Pravastatin	40 mg as a single dose	Fenofibrate 3 x 67 mg ² as a single dose	↓1%	↓2%
Fluvastatin	40 mg as a single dose	Fenofibrate 160 mg ¹ as a single dose	↓2%	↓10%
Anti-diabetic agents				
Glimepiride	1 mg as a single dose	Fenofibrate 145 mg ¹ once daily for 10 days	↑1%	↓1%
Metformin	850 mg three times daily for 10 days	Fenofibrate 54 mg ¹ three times daily for 10 days	↓9%	↓6%
Rosiglitazone	8 mg once daily for 5 days	Fenofibrate 145 mg ¹ once daily for 14 days	10%	†3%

¹Fenofibrate oral tablet ²Fenofibrate oral micronized capsule

Table 3. Effects of Fenofibrate Co-Administration on Systemic Exposure of Other

Dosage Regimen	Dosage Regimen of	Changes in Co-Administered			
of Fenofibrate Co-Administered Drug		Drug Exposure			
		Analyte		C _{max}	
Lipid-lowering agents					
Fenofibrate 160 mg ¹ once daily for 10 days			↓17%	0%	
Fenofibrate 3 x 67 mg ²	Pravastatin, 40 mg	Pravastatin	13%	13%	
as a single dose as a single dose Fenofibrate 160 mg ¹ Fluvastatin, 40 mg as a single dose as a single dose		3α-Hydroxyl- iso-pravastatin	↑26%	↑29%	
		(+)-3R, 5S- Fluvastatin	15%	16%	
Anti-diabetic agents					
Fenofibrate 145 mg ¹ once daily for 10 days	Glimepiride, 1 mg as a single dose	Glimepiride	135%	<u></u> †18%	
Fenofibrate 54 mg ¹ three times daily for 10 days	Metformin, 850 mg three times daily for 10 days	Metformin	†3%	↑6%	
Fenofibrate 145 mg ¹ once daily for 14 days	Rosiglitazone, 8 mg once daily for 5 days	Rosiglitazone	↑6%	↓1%	
¹ Fenofibrate oral tablet ² Fenofibrate oral micronized capsule					

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis and Mutagenesis and Impairment of Fertility

Two dietary carcinogenicity studies have been conducted in rats with fenofibrate. In the first 24-month study, Wistar rats were dosed with fenofibrate at 10, 45, and 200 mg/kg/day, approximately 0.3, 1, and 6 times the maximum recommended human dose (MRHD) of 300 mg fenofibrate daily, equivalent to 145 mg fenofibrate daily based on body surface area comparisons. At a dose of 200 mg/kg/day (at 6 times the MRHD), the incidence of liver carcinomas was significant increased in both sexes. A statistically significant increase in pancreatic adenomas and benign testicular interstitial cell tumors was observed at 6 times the MRHD) in males. In a second 24-month rat carcinogenicity study in a different strain of rats (Sprague-Dawley), doses of 10 and 60 mg/kg/day (0.3 and 2 times the MRHD) produced significant increases in the incidence of pancreatic acinomas in both sexes and increases in testicular interstitial cell tumors in males at 2 times the MRHD) model.

A 117-week carcinogenicity study was conducted in rats comparing three drugs: fenofibrate 10 and 60 mg/kg/day (0.3 and 2 times the MRHD, based on body surface area comparisons), clofibrate (400 mg/kg/day; 2 times the human dose), and gemfibrozil (250 mg/kg/day; 2 times the human dose, based on mg/m² surface area). Fenofibrate increased pancreatic acinar adenomas in both sexes. Clofibrate increased hepatocellular carcinoma and pancreatic acinar adenomas in males and hepatic neoplastic nodules in females. Gemfibrozil increased hepatic neoplastic nodules in males and females, while all three drugs increased testicular interstitial cell tumors in males.

In a 21-month study in CF-1 mice, fenofibrate 10, 45, and 200 mg/kg/day (approximately 0.2, 1, and 3 times the MRHD, based on body surface area comparisons) significantly increased the liver carcinomas in both sexes at 3 times the MRHD. In a second 18-month study at 10, 60, and 200 mg/kg/day, fenofibrate significantly increased the liver carcinomas in male mice and liver adenomas in female mice at 3 times the MRHD.

Electron microscopy studies have demonstrated peroxisomal proliferation following fenofibrate administration to the rat. An adequate study to test for peroxisome proliferation in humans has not been done, but changes in peroxisome morphology and numbers have been observed in humans after treatment with other members of the fibrate class when liver biopsies were compared before and after treatment in the same individual.

Fenofibrate has been demonstrated to be devoid of mutagenic potential in the following tests: Ames, mouse lymphoma, chromosomal aberration and unscheduled DNA synthesis in primary rat hepatocytes.

In fertility studies rats were given oral dietary doses of fenofibrate, males received 61 days prior to mating and females 15 days prior to mating through weaning which resulted in no adverse effect on fertility at doses up to 300 mg/kg/day (10 times the MRHD, based on body surface area comparisons).

14 CLINICAL STUDIES

14.1 Primary Hypercholesterolemia (Heterozygous Familial and Nonfamilial) and Mixed Dyslipidemia

The effects of fenofibrate at a dose equivalent to 145 mg fenofibrate tablets per day were assessed from four randomized, placebo-controlled, double-bind, parallel-group studies including patients with the following mean baseline lipid values: total-C 306.9 mg/dL; LDL-C 213.8 mg/dL; HDL-C 52.3 mg/dL; and triglycerides 191 mg/dL. Fenofibrate therapy lowered LDL-C, Total-C, and the LDL-C/HDL-C ratio. Fenofibrate therapy also lowered triglycerides and raised HDL-C (see Table 4).

Table 4. Mean Percent Change in Lipid Parameters at End of Treatment[†]

Treatment Group	Total-C	LDL-C	HDL-C	TG
Pooled Cohort				
Mean baseline lipid values (n = 646)	306.9 mg/dL	213.8 mg/dL	52.3 mg/dL	191 mg/dL
All FEN (n = 361)	-18.7%*	-20.6%*	+11%*	-28.9%*
Placebo (n = 285)	-0.4%	-2.2%	+0.7%	+7.7%
Baseline LDL-C > 160 mg/dL and TG < 150 mg/dL				
Mean baseline lipid values (n = 334)	307.7 mg/dL	227.7 mg/dL	58.1 mg/dL	101.7 mg/dL
All FEN (n = 193)	-22.4%*	-31.4%*	+9.8%*	-23.5%*
Placebo (n = 141)	+0.2%	-2.2%	+2.6%	+11.7%
Baseline LDL-C > 160 mg/dL and TG ≥ 150 mg/dL				
Mean baseline lipid values (n = 242)	312.8 mg/dL	219.8 mg/dL	46.7 mg/dL	231.9 mg/dL
All FEN (n = 126)	-16.8%*	-20.1%*	+14.6%*	-35.9%*
Placebo (n = 116)	-3%	-6.6%	+2.3%	+0.9%
†Duration of study treatment was 3 to 6 months. *p = < 0.05 vs. Placebo				

In a subset of the subjects, measurements of apo B were conducted. Fenofibrate treatment significantly reduced apo B from baseline to endpoint as compared with placebo (-25.1% vs. 2.4%, p < 0.0001, n = 213 and 143 respectively).

14.2 Severe Hypertriglyceridemia

The effects of fenofibrate on serum triglycerides were studied in two randomized, double-blind, placebo-controlled clinical trials of 147 hypertriglyceridemic patients. Patients were treated for eight weeks under protocols that differed only in that one entered patients with baseline TG levels of 500 to 1500 mg/dL, and the other TG levels of 350 to 500 mg/dL. In patients with hypertriglyceridemia and normal cholesterolemia with or without hyperchylomicronemia, treatment with fenofibrate at dosages equivalent to fenofibrate 145 mg per day decreased primarily very low density lipoprotein (VLDL) triglycerides and VLDL cholesterol. Treatment of patients with elevated triglycerides often results in an increase of LDL-C (see Table 5).

			e H	lypertriglyceridemia

Study 1	Plac	Placebo			Fend	Fenofibrate			
Baseline TG levels 350 to 499 mg/dL	N	Baseline (Mean)	Endpoint (Mean)	% Change (Mean)	N	Baseline (Mean)	Endpoint (Mean)	% Change (Mean)	
Triglycerides	28	449	450	-0.5	27	432	223	-46.2*	
VLDL Triglycerides	19	367	350	2.7	19	350	178	-44.1	
Total Cholesterol	28	255	261	2.8	27	252	227	-9.1	
HDL Cholesterol	28	35	36	4	27	34	40	19.6*	
LDL Cholesterol	28	120	129	12	27	128	137	14.5	
VLDL Cholesterol	27	99	99	5.8	27	92	46	-44.7	
Study 2	Plac	Placebo			Fenofibrate				
Baseline TG levels 500 to 1500 mg/dL	N	Baseline (Mean)	Endpoint (Mean)	% Change (Mean)	N	Baseline (Mean)	Endpoint (Mean)	% Change (Mean)	
Triglycerides	44	710	750	7.2	48	726	308	-54.5*	
VLDL Triglycerides	29	537	571	18.7	33	543	205	-50.6	
Total Cholesterol	44	272	271	0.4	48	261	223	-13.8	
HDL Cholesterol	44	27	28	5	48	30	36	22.9*	
LDL Cholesterol	42	100	90	-4.2	45	103	131	45	
VLDL Cholesterol	42	137	142	11	45	126	54	-49.4	
* = p < 0.05 vs. Placebo									

The effect of fenofibrate on cardiovascular morbidity and mortality has not beer determined.

16 HOW SUPPLIED/STORAGE AND HANDLING

Fenofibrate tablets, USP are available in two strengths: 48 mg white to off-white, oval-shaped, biconvex, film-coated tablets debossed with 'RH37' on one side and other side plain and are available as follows:

NDC 63304-448-30	Bottles of 30
NDC 63304-448-90	Bottles of 90

NDC 63304-448-	-05 Botti	es of 500			
145 mg white to	off-white, oval-shaped	l, biconvex,	film-coated	tablets debossed	with

'RH38' on one side and other side plain and are available as follows: NDC 63304-449-30 Bottles of 30

torano						
DC 63304-449-05	Bottles of 500					
DC 63304-449-90	Bottles of 90					
00 00004 440 00	Dottios of 00					

Store at 20° - 25° C (68° - 77° F) [See USP Controlled Room Temperature]. Keep out of the reach of children. Protect from moisture.

17 PATIENT COUNSELING INFORMATION

tablets, USP

taking fenofibrate tablets, USP.

after the final dose.

Sun Pharmaceutical Industries Ltd. Mohali, INDIA

Sun Pharmaceutical Industries, Inc. Cranbury, NJ 08512

Manufactured by:

Distributed by:

July 2021

Patients should be advised: • of the potential benefits and risks of fenofibrate tablets, USP.

prescribed dose, swallowing each tablet whole. to return to their physician's office for routine monitoring.

- not to use fenofibrate tablets, USP if there is a known hypersensitivity to fenofibrate or fenofibric acid.
 of medications that should not be taken in combination with fenofibrate tablets,
- of medications that should not be taken in combination with fenofibrate tablets, USP.
 that if they are taking coumarin anticoagulants, fenofibrate tablets. USP may

increase their anti-cogulant effect, and increased monitoring may be necessary, to continue to follow an appropriate lipid-modifying diet while taking fenofibrate

to take fenofibrate tablets. USP once daily, without regard to food, at the

to inform their physician of all medications, supplements, and herbal preparations

they are taking and any charge to their medical condition. Patients should also be advised to inform their physicians prescribing a new medication that they are

to inform their physician of symptoms of liver injury (e.g., jaundice, abdominal pain, nausea, malaise, dark urine, abnormal stool, pruritus); any muscle pain, tenderness, or weakness; or any other new symptoms.

not to breastfeed during treatment with fenofibrate tablets, USP and for 5 days

FDA-08

486

5221