



Dispense with Patient Information available at: <https://www.sunpharma.com/usa/products>

PATIENT INFORMATION Rosuvastatin (roe-SOO-va-STAT-in) Tablets, USP
Read this Patient Information carefully before you start taking rosuvastatin tablets and each time you get a refill. If you have any questions about rosuvastatin tablets, ask your doctor. Only your doctor can determine if rosuvastatin tablets are right for you.
What is rosuvastatin tablets? Rosuvastatin tablets are prescription medicine that contains a cholesterol-lowering medicine called rosuvastatin. <ul style="list-style-type: none">• Rosuvastatin tablets are used to:<ul style="list-style-type: none">○ reduce the risk of stroke, heart attack, and the need for procedures to improve blood flow to the heart called arterial revascularization in adults who do not have known heart disease but do have certain additional risk factors.• Rosuvastatin tablets are used along with diet to:<ul style="list-style-type: none">○ lower the level of low-density lipoprotein (LDL) cholesterol or “bad” cholesterol in adults with primary hyperlipidemia.○ slow the buildup of fatty deposits (plaque) in the walls of blood vessels.○ treat adults and children 8 years of age and older with high blood cholesterol due to heterozygous familial hypercholesterolemia (an inherited condition that causes high levels of LDL).○ treat adults with a type of high cholesterol called primary dysbetalipoproteinemia (type III hyperlipoproteinemia).○ lower the level of fat in your blood (triglycerides) in adults with hypertriglyceridemia. <p>The safety and effectiveness of rosuvastatin tablets has not been established in children younger than 8 years of age with heterozygous familial hypercholesterolemia or in children with other types of hyperlipidemias (other than HeFH or HoFH).</p> Pediatric use information for patients 7 to 17 years of age is approved for AstraZeneca's CRESTOR (rosuvastatin) tablets. However, due to AstraZeneca's marketing exclusivity rights, this drug product is not labeled with that pediatric information.
Who should not take rosuvastatin tablets? Do not take rosuvastatin tablets if you: <ul style="list-style-type: none">• have liver problems.• are allergic to rosuvastatin or any of the ingredients in rosuvastatin tablets. See the end of this leaflet for a complete list of ingredients in rosuvastatin tablets. Symptoms of allergic reactions include rash, itching, hives, and swelling.

What should I tell my doctor before and while taking rosuvastatin tablets? Tell your doctor if you: <ul style="list-style-type: none">• have unexplained muscle aches or weakness.• have or have had kidney problems.• have or have had liver problems.• drink more than 2 glasses of alcohol daily.• have thyroid problems.• are 65 years of age or older.• are of Asian descent.• are pregnant or think you may be pregnant, or are planning to become pregnant. If you become pregnant while taking rosuvastatin tablets, call your healthcare provider right away to discuss your rosuvastatin tablets treatment.• are breastfeeding. Rosuvastatin tablets can pass into your breast milk. Breastfeeding is not recommended while taking rosuvastatin tablets. Tell your doctor about all the medicines you take , including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking rosuvastatin tablets with certain other medicines may affect each other causing side effects. Rosuvastatin tablets may affect the way other medicines work, and other medicines may affect how rosuvastatin tablets work. Especially tell your doctor if you take: <ul style="list-style-type: none">• cyclosporine (a medicine for your immune system)• gemfibrozil (a fibric acid medicine for lowering cholesterol)• fostamatinib (a medicine used to treat low platelet counts)• febuxostat (a medicine used to treat and prevent high blood levels of uric acid)• teriflunomide (a medicine used to treat relapsing remitting multiple sclerosis)• capmatinib (a medicine for the treatment of non-small cell lung cancer)• tafamidis (used to treat cardiomyopathy [enlarged and thickened heart muscle])• darolutamide (a medicine for the treatment of prostate cancer)• regorafenib (a medicine used to treat cancer of the colon and rectum)• anti-viral medicines including certain HIV or hepatitis C virus drugs such as:<ul style="list-style-type: none">○ lopinavir, ritonavir, fosamprenavir, tipranavir, atazanavir, simeprevir○ combination of<ul style="list-style-type: none">■ sofosbuvir/velpatasvir/voxilaprevir■ dasabuvir/ombitasvir/paritaprevir/ritonavir■ elbasvir/grazoprevir■ sofosbuvir/velpatasvir■ glecaprevir/pibrentasvir and○ all other combinations with ledipasvir including ledipasvir/sofosbuvir• certain anti-fungal medicines (such as itraconazole, ketoconazole and fluconazole)• coumarin anticoagulants (medicines that prevent blood clots, such as warfarin)• niacin or nicotinic acid• fibric acid derivatives (such as fenofibrate)• colchicine (a medicine used to treat gout) Ask your doctor or pharmacist for a list of these medicines if you are not sure. Know the medicines you take. Keep a list of them to show your doctor and pharmacist when you get new medicine.
--

How should I take rosuvastatin tablets? <ul style="list-style-type: none">• Take rosuvastatin tablets exactly as your doctor tells you to take it.• Take rosuvastatin tablets, by mouth, 1 time each day. Swallow the tablet whole.• Rosuvastatin tablets can be taken at any time of day, with or without food.• Do not change your dose or stop rosuvastatin tablets without talking to your doctor, even if you are feeling well.• Your doctor may do blood tests to check your cholesterol levels before and during your treatment with rosuvastatin tablets. Your doctor may change your dose of rosuvastatin tablets if needed.• Your doctor may start you on a cholesterol lowering diet before giving you rosuvastatin tablets. Stay on this diet when you take rosuvastatin tablets.• Wait at least 2 hours after taking rosuvastatin tablets to take an antacid that contains a combination of aluminum and magnesium hydroxide.• If you miss a dose of rosuvastatin tablets, take your next dose at your normal scheduled time. Do not take an extra dose of rosuvastatin tablets.• If you take too many rosuvastatin tablets or overdose, call your doctor or go to the nearest hospital emergency room right away.
What are the possible side effects of rosuvastatin tablets? Rosuvastatin tablets may cause serious side effects, including: <ul style="list-style-type: none">• Muscle pain, tenderness and weakness (myopathy). Muscle problems, including muscle breakdown, can be serious in some people and rarely cause kidney damage that can lead to death. Tell your doctor right away if:<ul style="list-style-type: none">○ you have unexplained muscle pain, tenderness, or weakness, especially if you have a fever or feel more tired than usual, while you take rosuvastatin tablets.○ you have muscle problems that do not go away even after your doctor has told you to stop taking rosuvastatin tablets. Your doctor may do further tests to diagnose the cause of your muscle problems. Your chances of getting muscle problems are higher if you: <ul style="list-style-type: none">○ are taking certain other medicines while you take rosuvastatin tablets○ are 65 years of age or older○ have thyroid problems (hypothyroidism) that are not controlled○ have kidney problems○ are taking higher doses of rosuvastatin tablets <ul style="list-style-type: none">• Liver problems. Your doctor should do blood tests to check your liver before you start taking rosuvastatin tablets and if you have symptoms of liver problems while you take rosuvastatin tablets. Call your doctor right away if you have any of the following symptoms of liver problems:<ul style="list-style-type: none">○ feel unusually tired or weak○ loss of appetite○ upper belly pain○ dark urine○ yellowing of your skin or the whites of your eyes• Protein and blood in the urine. Rosuvastatin tablets may cause you to have protein and blood in your urine. If you develop protein or blood in your urine, your doctor may decrease your dose of rosuvastatin tablets.• Increase in blood sugar (glucose) levels. Rosuvastatin tablets may cause an increase in your blood sugar levels. The most common side effects may include headache, muscle aches and pains, abdominal pain, weakness, and nausea.

Tell your doctor if you have any side effect that bothers you or that does not go away.
For more information, ask your doctor or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to
FDA at 1-800-FDA-1088.

How should I store rosuvastatin tablets?

- Store rosuvastatin tablets at 20° to 25°C (68° to 77°F) and in a dry place.
- Safely throw away medicine that is out of date or no longer needed.
- Rosuvastatin tablets come in a child-resistant package.

Keep rosuvastatin tablets and all medicines out of the reach of children.

General Information about the safe and effective use of rosuvastatin tablets

Medicines are sometimes prescribed for purposes other than those listed in a Patient
Information leaflet. Do not use rosuvastatin tablets for a condition for which it was
not prescribed. Do not give rosuvastatin tablets to other people, even if they have the
same medical condition you have. It may harm them.

You can ask your pharmacist or doctor for information about rosuvastatin tablets
that is written for health professionals.

What are the Ingredients in Rosuvastatin tablets?

Active Ingredient: rosuvastatin as rosuvastatin calcium

Inactive Ingredients: colloidal silicon dioxide, hypromellose, magnesium stearate,
microcrystalline cellulose, polyethylene glycol, pregelatinized maize starch, talc,
titanium dioxide, iron oxide yellow (5 mg), iron oxide red (10 mg, 20 mg, 40 mg), and
FD&C Red # 40 aluminum lake (10 mg, 20 mg, 40 mg).

Distributed by:

Sun Pharmaceutical Industries, Inc.

Cranbury, NJ 08512



Manufactured by:

Sun Pharmaceutical Industries Ltd.

Survey No. 259/15, Dadra-396 191,
(U.T. of D & NH), India.

5239355
Revised 02/2023

For more information, call 1-800-818-4555.

This Patient Information has been approved by the U.S. Food and Drug
Administration