

Dispense with Medication Guide available at: https://www.sunpharma.com/usa/products

Medication Guide

Venlafaxine Hydrochloride Extended-Release Tablets

(ven-luh-fak-seen HYE-droe-KLOR-ide)

Antidepressant Medicines, Depression and Other Serious Mental Illnesses, and Suicidal **Thoughts or Actions**

member's antidepressant medicine. This Medication Guide is only about the risk of suicidal thoughts and actions with antidepressant medicines. Talk to your, or your family member's, healthcare provider about:

Read the Medication Guide that comes with your or your family

- all risks and benefits of treatment with antidepressant medicines - all treatment choices for depression or other serious mental
 - illness

Who should not take venlafaxine hydrochloride extendedrelease tablets?

- take a monoamine oxidase inhibitor (MAOI). Ask your healthcare provider or pharmacist if you are not sure if you take an MAOI, including the antibiotic linezolid.
- Do not take an MAOI within 7 days of stopping venlafaxine hydrochloride extended-release tablets unless directed to do so by your physician.
- Do not start venlafaxine hydrochloride extended-release tablets if you stopped taking an MAOI in the last 2 weeks unless directed to do so by your physician.

What is the most important information I should know about antidepressant medicines, depression and other serious mental illnesses, and suicidal thoughts or actions? 1. Antidepressant medicines may increase suicidal thoughts or actions in some children, teenagers, and young adults within

the first few months of treatment. 2. Depression and other serious mental illnesses are the most important causes of suicidal thoughts and actions. Some people may have a particularly high risk of having suicidal

thoughts or actions. These include people who have (or have a family history of) bipolar illness (also called manic-depressive illness) or suicidal thoughts or actions.

3. How can I watch for and try to prevent suicidal thoughts and actions in myself or a family member?

when an antidepressant medicine is started or when the dose is

Pay close attention to any changes, especially sudden changes. in mood, behaviors, thoughts, or feelings. This is very important

changed. - Call the healthcare provider right away to report new or sudden

changes in mood, behavior, thoughts, or feelings. - Keep all follow-up visits with the healthcare provider as

scheduled. Call the healthcare provider between visits as needed, especially if you have concerns about symptoms.

Call a healthcare provider right away if you or your family

member has any of the following symptoms, especially if they are new, worse, or worry you: - thoughts about suicide or dving

- attempts to commit suicide - new or worse depression

- new or worse anxiety - feeling very agitated or restless - panic attacks

- trouble sleeping (insomnia) - new or worse irritability

around eve

- acting aggressive, being angry, or violent - acting on dangerous impulses

- an extreme increase in activity and talking (mania)

- other unusual changes in behavior or mood

- Visual Problems: Eye pain, change in vision, swelling or redness

What else do I need to know about antidepressant medicines? - Never stop an antidepressant medicine without first talking to a healthcare provider. Stopping an antidepressant medicine suddenly can cause other symptoms.

- Visual Problems. Only some people are at risk for these problems. You may want to undergo an eye examination to see if you are at risk and receive preventative treatment if you are. - Antidepressants are medicines used to treat depression and

other illnesses. It is important to discuss all the risks of treating depression and also the risks of not treating it. Patients and their families or other caregivers should discuss all treatment choices with the healthcare provider, not just the use of antidepressants.

- Antidepressant medicines have other side effects. Talk to the healthcare provider about the side effects of the medicine prescribed for you or your family member. Antidepressant medicines can interact with other medicines.
- Know all of the medicines that you or your family member takes. Keep a list of all medicines to show the healthcare provider. Do

not start new medicines without first checking with your healthcare provider.

Especially tell your healthcare provider if you take:

o medicines to treat migraine headaches known as triptans

o tricyclic antidepressants o lithium o tramadol, fentanyl, meperidine, methadone, or other opioids

o tryptophan o buspirone o amphetamines

o St. John's Wort o phentermine

sexual problems.

o other medicines containing desvenlafaxine or venlafaxine o medicines that can affect blood clotting such as aspirin.

nonsteroidal anti-inflammatory drugs (NSAIDs), warfarin - Not all antidepressant medicines prescribed for children are FDA approved for use in children. Talk to your child's

healthcare provider for more information. Sexual problems (dysfunction). Taking serotonin and norepinephrine reuptake inhibitors (SNRIs), including venlafaxine hydrochloride extended-release tablets, may cause

Symptoms in males may include:

o Delayed ejaculation or inability to have an ejaculation

o Decreased sex drive

o Problems getting or keeping an erection

Symptoms in females may include:

o Decreased sex drive

o Delayed orgasm or inability to have an orgasm

Talk to your healthcare provider if you develop any changes in your sexual function or if you have any questions or concerns about sexual problems during treatment with venlafaxine hydrochloride extended-release tablets. There may be treatments your healthcare provider can suggest.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088

This Medication Guide has been approved by the U.S. Food and Drug Administration for all antidepressants.

Distributed by: Sun Pharmaceutical Industries, Inc. Cranbury, NJ 08512

Manufactured by: Sun Pharmaceutical Industries Ltd.

Halol-Baroda Highway, PHARMA Halol-389 350, Gujarat, India.