

Ranbaxy Pharmaceuticals (Pty) Ltd

GEMVALAZ 5/160 and 10/160

Each film-coated tablet contains 6,93 mg and 13,87 mg amlodipine besylate (equivalent to 5 mg and 10 mg amlodipine base) respectively and 160 mg valsartan

PATIENT INFORMATION LEAFLET

SCHEDULING STATUS: S3

GEMVALAZ 5/160 mg tablet (5 mg amlodipine and 160 mg valsartan)

GEMVALAZ 10/160 mg tablet (10 mg amlodipine and 160 mg valsartan)

Sugar free

Read all of this leaflet carefully before you start taking GEMVALAZ

- Keep this leaflet. You may need to read it again.
- If you have further questions, please ask your doctor, pharmacist, nurse or other healthcare provider.
- GEMVALAZ has been prescribed for you personally and you should not share your medicine with other people. It may harm them, even if their symptoms are the same as yours.

What is in this leaflet

1. What GEMVALAZ is and what it is used for
2. What you need to know before you take GEMVALAZ
3. How to take GEMVALAZ
4. Possible side effects
5. How to store GEMVALAZ
6. Contents of the pack and other information

1. What GEMVALAZ is and what it is used for

GEMVALAZ tablets contain two substances called amlodipine and valsartan. Both of these substances help to control high blood pressure.

Signed: *Carlye Aizer*

Ranbaxy Pharmaceuticals (Pty) Ltd

GEMVALAZ 5/160 and 10/160

Each film-coated tablet contains 6,93 mg and 13,87 mg amlodipine besylate (equivalent to 5 mg and 10 mg amlodipine base) respectively and 160 mg valsartan

Amlodipine belongs to a group of substances called “calcium channel blockers”. Amlodipine stops calcium from moving into the blood vessel wall which stops the blood vessels from tightening.

Valsartan belongs to a group of substances called “angiotensin-II receptor antagonists”. Angiotensin II is produced by the body and makes the blood vessels tighten, thus increasing the blood pressure. Valsartan works by blocking the effect of angiotensin II.

This means that both of these substances help to stop the blood vessels tightening. As a result, the blood vessels relax and blood pressure is lowered.

GEMVALAZ is used to treat high blood pressure in adults whose blood pressure is not controlled enough with either amlodipine or valsartan on its own.

2. What you need to know before you take GEMVALAZ

Do not take GEMVALAZ:

- If you are hypersensitive (allergic) to amlodipine or any other calcium channel blockers, valsartan or to any of the other ingredients of GEMVALAZ (listed in section 6).
- If you have experienced swelling, particularly of the face and throat, while taking other medicines (including angiotensin converting enzyme inhibitors). If you get these symptoms, stop taking GEMVALAZ and contact your doctor straight away. You should never take GEMVALAZ again.
- If you have recurrent episodes of severe swelling without a known cause.
- Your doctor has diagnosed that you are suffering from a narrowing of valves in your heart (called aortic or mitral stenosis), or abnormally increased thickness of your heart muscle with narrowing (called obstructive hypertrophic cardiomyopathy)

Ranbaxy Pharmaceuticals (Pty) Ltd

GEMVALAZ 5/160 and 10/160

Each film-coated tablet contains 6,93 mg and 13,87 mg amlodipine besylate (equivalent to 5 mg and 10 mg amlodipine base) respectively and 160 mg valsartan

- If you have received a kidney transplant or if you had been diagnosed to suffer from a narrowing of your kidney artery or if you are suffering from severe kidney problems
 - If you are taking other medicine or substances, which increase the potassium levels in your blood (such as certain types of diuretics, potassium supplements, etc.)
 - If you have porphyria
 - If you are taking a medicine called lithium
 - If you are treated with a blood pressure lowering medicine containing aliskiren.
 - If you have severe liver impairment
 - If you have severe low blood pressure (hypotension)
 - If you have cardiogenic shock (a condition where your heart is unable to supply enough blood to the body)
 - If you suffer from heart failure after a heart attack
 - If you are pregnant or breastfeeding.

Warnings and precautions

Take special care with GEMVALAZ:

- if you are taking aliskiren (see “Do not take”)
- if you have been sick (vomiting or diarrhoea)
- if you have had a kidney transplant or if you had been told that you have a narrowing of your kidney arteries
- if you have liver or kidney problems
- if you have a condition affecting the renal glands called “primary hyperaldosteronism”.
- if you have experienced swelling, particularly of the face and throat, while taking other medicines (including angiotensin converting enzyme inhibitors). If you get these symptoms, stop taking GEMVALAZ and contact your doctor straight away. You should never take GEMVALAZ again.

Signed: *Carlysa Aizer*

Ranbaxy Pharmaceuticals (Pty) Ltd

GEMVALAZ 5/160 and 10/160

Each film-coated tablet contains 6,93 mg and 13,87 mg amlodipine besylate (equivalent to 5 mg and 10 mg amlodipine base) respectively and 160 mg valsartan

- if you have had heart failure or have experienced a heart attack. Follow your doctor's instructions for the starting dose carefully. Your doctor may also check your kidney function.

Contact your doctor to re-evaluate your treatment if you are treated with ACE

Inhibitors/Angiotensin receptor blockers together with a fluoroquinolone antibiotic.

Your doctor may check your kidney function, blood pressure, and the amount of electrolytes (e.g. potassium) in your blood at regular intervals.

Children and adolescents

The use of GEMVALAZ in children and adolescents is not recommended (aged below 18 years old).

Other medicines and GEMVALAZ

Always tell your healthcare provider if you are taking any other medicine. (This includes all complementary or traditional medicines.)

Tell your doctor if you are taking:

- ACE inhibitors or aliskiren (medicines used to treat high blood pressure) (see also information under the headings "Do not take GEMVALAZ")
- diuretics (a type of medicine also called "water tablets" which increases the amount of urine you produce)
- potassium-sparing diuretics, potassium supplements, salt substitutes containing potassium and other substances that may increase potassium levels
- nitroglycerin and other nitrates, or other substances called "vasodilators"
- medicines used for HIV/AIDS (e.g. ritonavir, indinavir, nelfinavir)

Signed: *Carlye Aizer*

Ranbaxy Pharmaceuticals (Pty) Ltd**GEMVALAZ 5/160 and 10/160**

Each film-coated tablet contains 6,93 mg and 13,87 mg amlodipine besylate (equivalent to 5 mg and 10 mg amlodipine base) respectively and 160 mg valsartan

- medicines used to treat fungal infections (e.g. ketoconazole, itraconazole)
- clarithromycin (for infections caused by bacteria)
- verapamil, diltiazem (heart medicines)
- medicines used to treat bacterial infections (such as rifampicin, erythromycin, talithromycin)
- anticonvulsant medicines (e.g. carbamazepine, phenobarbitone, phenytoin, fosphenytoin, primidone)
- St. John's wort
- simvastatin (a medicine used to control high cholesterol levels)
- dantrolene (infusion for severe body temperature abnormalities)
- tacrolimus (used to control your body's immune response, enabling your body to accept the transplanted organ).
- medicines used to protect against transplant rejection (ciclosporin)
- lithium (a medicine used to treat some types of depression) (see also information under the headings "Do not take GEMVALAZ")
- certain types of painkillers called non-steroidal anti-inflammatory medicines (NSAIDs) or selective cyclooxygenase-2 inhibitors (COX-2 inhibitors). Your doctor may also check your kidney function

GEMVALAZ with food and drink

Grapefruit and grapefruit juice should not be consumed by people who are taking GEMVALAZ. This is because grapefruit and grapefruit juice can lead to an increase in the blood levels of the active substance amlodipine, which can cause an unpredictable increase in the blood pressure lowering effect of GEMVALAZ

Pregnancy and breastfeeding

Signed: *Carlye Aizer*

Ranbaxy Pharmaceuticals (Pty) Ltd

GEMVALAZ 5/160 and 10/160

Each film-coated tablet contains 6,93 mg and 13,87 mg amlodipine besylate (equivalent to 5 mg and 10 mg amlodipine base) respectively and 160 mg valsartan

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, please consult your doctor, pharmacist or other healthcare provider for advice before taking GEMVALAZ.

Safety of GEMVALAZ in pregnant and breastfeeding women has not been established.

You should not use GEMVALAZ if you are pregnant, trying to become pregnant or breastfeeding your baby.

If you are a woman of childbearing age, you should ensure that you use an effective contraception.

Driving and using machines

This medicine may make you feel dizzy. This can affect how well you can concentrate.

It is not always possible to predict to what extent GEMVALAZ may interfere with your daily activities. You should ensure that you do not engage in driving a vehicle or operate machines until you are aware of the measure to which GEMVALAZ affects you.

GEMVALAZ is sugar free.

3. How to take GEMVALAZ

Do not share medicines prescribed for you with any other person.

Always take GEMVALAZ exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

The usual dose of GEMVALAZ is one tablet per day.

Signed: *Carlyne Aizer*

Ranbaxy Pharmaceuticals (Pty) Ltd

GEMVALAZ 5/160 and 10/160

Each film-coated tablet contains 6,93 mg and 13,87 mg amlodipine besylate (equivalent to 5 mg and 10 mg amlodipine base) respectively and 160 mg valsartan

- It is preferable to take your medicine at the same time each day.
- Swallow the tablets with a glass of water.
- You can take GEMVALAZ with or without food. Do not take GEMVALAZ with grapefruit or grapefruit juice.

Depending on how you respond to the treatment, your doctor may suggest a higher or lower dose.

Do not exceed the prescribed dose.

GEMVALAZ and older people (age 65 years or over)

Your doctor should exercise caution when increasing your dose.

If you take more GEMVALAZ than you should

In the event of overdose, consult your doctor or pharmacist. If neither is available, contact the nearest hospital or poison centre. You may experience dizziness, difficulty in breathing and a faster heartbeat.

If you forget to take GEMVALAZ

Do not take a double dose to make up for forgotten individual doses.

If you stop taking GEMVALAZ

Stopping your treatment with GEMVALAZ may cause your disease to get worse. Do not stop taking your medicine unless your doctor tells you to.

Signed: *Carlye Aizer*

Ranbaxy Pharmaceuticals (Pty) Ltd

GEMVALAZ 5/160 and 10/160

Each film-coated tablet contains 6,93 mg and 13,87 mg amlodipine besylate (equivalent to 5 mg and 10 mg amlodipine base) respectively and 160 mg valsartan

4. Possible side effects

GEMVALAZ can have side effects.

Not all side effects reported for GEMVALAZ are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking GEMVALAZ, please consult your healthcare provider for advice.

If any of the following symptoms occur, stop taking the GEMVALAZ and tell your doctor immediately or go to the casualty department at your nearest hospital:

- swelling of the hands, feet, ankles, face, lips and mouth or throat, which may cause difficulty in swallowing or breathing,
- rash or itching,
- fainting.

These are all very serious side effects. If you have them, you may have had a serious allergic reaction to GEMVALAZ. You may need urgent medical attention or hospitalisation.

Tell your doctor immediately or go to the casualty department at your nearest hospital if you notice any of the following:

- sudden wheeziness, chest pain, shortness of breath or difficulty in breathing
- severe skin reactions including intense skin rash, hives, reddening of the skin over your whole body, severe itching, blistering, peeling and swelling of the skin, inflammation of the mucous membranes (Stevens-Johnson Syndrome, toxic epidermal necrolysis) or other allergic reactions
- inflamed pancreas, which may cause severe abdominal and back pain accompanied with feeling of being very unwell
- heart attack, abnormal heartbeat

Signed: *Carlye Aizer*

Ranbaxy Pharmaceuticals (Pty) Ltd

GEMVALAZ 5/160 and 10/160

Each film-coated tablet contains 6,93 mg and 13,87 mg amlodipine besylate (equivalent to 5 mg and 10 mg amlodipine base) respectively and 160 mg valsartan

- upper airway swelling

These are all serious side effects. You may need urgent medical attention. Tell your doctor if you notice any of the following:

Frequent side effects

- influenza (flu); blocked nose, sore throat and discomfort when swallowing
- low potassium levels
- headache
- asthenia (weakness)
- dizziness
- stomach pain
- feeling sick (nausea)
- tiredness
- facial swelling
- redness and warm feeling of the face and/or neck
- swelling in the body caused by excess fluid. It often affects the lower body, such as the legs, feet, and ankle but it can occur anywhere

Less frequent side effects

- loss of appetite
- high calcium, sugar, cholesterol or uric acid levels in the blood
- low levels of sodium in the blood
- feeling anxious
- abnormal coordination
- drowsiness, tingling or numbness of the hands or feet
- vision disturbance and impairment
- ringing in the ears, head spinning

Signed: *Carlye Aizer*

Ranbaxy Pharmaceuticals (Pty) Ltd

GEMVALAZ 5/160 and 10/160

Each film-coated tablet contains 6,93 mg and 13,87 mg amlodipine besylate (equivalent to 5 mg and 10 mg amlodipine base) respectively and 160 mg valsartan

- fast heartbeat including palpitations
- low blood pressure and light-headedness when standing up
- cough, sneezing/runny nose caused by inflammation of the lining of the nose (rhinitis)
- throat pain
- constipation, change of bowel habit,
- diarrhoea, dry mouth
- nausea, vomiting, indigestion
- excessive sweating
- redness of the skin
- skin rash all over your body
- joint swelling
- back pain; pain in joints, muscle spasm, muscle cramps
- sensation of heaviness
- passing more urine than normal or feeling more of an urge to pass urine
- inability to get or maintain an erection
- a low level of white blood cells in the blood
- low blood platelet count
- excess sugar in blood (hyperglycaemia)
- mood changes, depression, confusion
- taste abnormalities, loss of pain sensation, abnormal sensation, typically tingling or pricking ('pins and needles'), high level of muscle tone
- trembling, sleeplessness, partial or total loss of sensation in a part of your body
- an inflammation of the blood vessels that causes changes in the blood vessel walls
- swelling of the gums

Signed: *Carlye Aizer*

Ranbaxy Pharmaceuticals (Pty) Ltd

GEMVALAZ 5/160 and 10/160

Each film-coated tablet contains 6,93 mg and 13,87 mg amlodipine besylate (equivalent to 5 mg and 10 mg amlodipine base) respectively and 160 mg valsartan

- abdominal bloating (gastritis), increased liver enzyme increase which may have an effect on some medical tests
- abnormal liver function, inflammation of the liver (hepatitis), yellowing of the skin (jaundice)
- hair loss
- skin discolouration
- skin sensitivity to light, purple spots under the skin, hives
- discomfort or enlargement of the breasts in men
- lack of energy
- chest pain, feeling unwell
- pain
- weight increase or decrease

If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

Reporting of side effects

If you get side effects, talk to your doctor, pharmacist or nurse. You can also report side effects to SAHPRA via the “**6.04 Adverse Drug Reaction Reporting Form**”, found online under SAHPRA’s publications: <https://www.sahpra.org.za/Publications/Index/8>. By reporting side effects, you can help provide more information on the safety of GEMVALAZ.

5. How to store GEMVALAZ

STORE ALL MEDICINES OUT OF REACH OF CHILDREN.

Store at or below 25 °C in original package.

Protect from moisture.

Do not use this medicine after the expiry date which is stated on the carton and blister.

The expiry date refers to the last day of that month.

Signed: *Carlye Aizer*

Ranbaxy Pharmaceuticals (Pty) Ltd

GEMVALAZ 5/160 and 10/160

Each film-coated tablet contains 6,93 mg and 13,87 mg amlodipine besylate (equivalent to 5 mg and 10 mg amlodipine base) respectively and 160 mg valsartan

Keep the blisters in the outer carton in order to protect from moisture.

Do not throw away any medicines via wastewater or household waste.

Ask your pharmacist how to throw away medicines you no longer use.

These measures will help protect the environment

6. Contents of the pack and other information

What GEMVALAZ contains

The active substances are a combination of amlodipine besilate and valsartan.

GEMVALAZ 5/160 mg (Film-coated tablets)

Each film-coated tablet contains:

Valsartan 160 mg

Amlodipine besilate 6,93 mg equivalent to amlodipine 5 mg.

GEMVALAZ 10/160 mg (Film-coated tablets)

Valsartan 160 mg

Amlodipine besilate 13,87 mg equivalent to Amlodipine 10 mg.

The other ingredients are: Crospovidone, silica colloidal anhydrous, magnesium stearate, starch pre-gelatinized (starch 1500), microcrystalline cellulose, iron oxide yellow, talc, purified water, Opadry (03F82330) yellow, and Opadry (03F83059) orange.

What GEMVALAZ looks like and contents of the pack

GEMVALAZ 5/160 mg: Dark yellow colored, oval film coated tablet. debossed on one side with AV2 and plain on the other side.

GEMVALAZ 10/160 mg: Beige colored, oval film coated tablet debossed on one side with AV3 and plain on the other side.

Holder of Certificate of Registration

Signed: *Carlysa Aizer*

Ranbaxy Pharmaceuticals (Pty) Ltd

GEMVALAZ 5/160 and 10/160

Each film-coated tablet contains 6,93 mg and 13,87 mg amlodipine besylate (equivalent to 5 mg and 10 mg amlodipine base) respectively and 160 mg valsartan

RANBAXY PHARMACEUTICALS (PTY) LTD

14 Lautre Road,

Stormill, Ext.1, Roodepoort,

Johannesburg, 1724

South Africa

This leaflet was last revised in

08 November 2023

Registration number

Gemvalaz 5/160: 49/7.1.3/0167

Gemvalaz 10/160: 49/7.1.3/0167

Signed: *Carlye Aizer*